

European Public Health Association (EUPHA)

**Research workshop**, arranged by the EUPHA Section of Social security and health, in Lisbon 2008

**Title: Why do women have higher sickness absence than men? A review of suggested explanations and their connection to gender theories of health and social security.**

Chairpersons: Gunnel Hensing, Department of Public Health and Community Medicine, University of Gothenburg

Organiser: EUPHA section on Social Security and Health, Prof. Gunnel Hensing (PhD), Doctoral students Kristina Holmgren, Helena Rohdén and Carin Staland Nyman, Department of Public Health and Community Medicine, University of Gothenburg, Sweden

**Workshop abstract:**

The workshop includes a short introduction and four presentations with discussion.

Women have a higher incidence of sickness absence than men. This has repeatedly been shown in studies and statistics over the last 50 years. Still there is no coherent explanation or theory that can contribute to the development of interventions at work or in society to reduce this gender inequality in public health. Several explanations have been put forward. They can be grouped into five main clusters: burden of illness; social inequalities at work and in the labour market; the construction of the sickness insurance; treatment and rehabilitation in the health services and at work; and women's double work. An obvious but not much evolved explanation is that women's higher sickness absence is a mere reflection of sex differences in health and illness or of women's overall higher burden of illness. Several researchers have put forward that sex differences in sickness absence in reality are related to socioeconomic inequalities rather than gendered inequalities based mainly in detrimental exposures at work. The construction of the sickness insurance is important for the levels of sickness absence in different societies but few studies have analysed sickness insurances from a gender perspective. A growing number of studies focus on how women and men are treated in the health service, and to what extent men and women are provided with similar rehabilitation measures in the health services and at work. Finally, there are some studies focusing on women's higher responsibility for unpaid work and family obligations and the relation to sickness absence. In this workshop these perspectives, except for the construction of the sickness insurance, will be described in more detail, and their connection to gender theories on health and social security will be explored to increase the understanding of how a gender perspective can be developed in sickness absence research, but also to provide new angles to public health interventions.

Abstract 1 title: Health problems and sickness absence in women – what are the associations?

Abstract 1 presenting authors: Gunnel Hensing

Abstract 1 authors: Hensing G

Affiliations 1: Department of Public Health and Community Medicine, University of Gothenburg, Gothenburg, Sweden

Abstract 1: Issue

Why do women have higher levels of sickness absence in most societies and in most diagnostic groups compared to men?

Description of the problem

Whether sickness absence is a measure of ill health, of motivation to work, a reflection of the labour market situation or of the generosity of the social insurance system has been an issue for researchers of different disciplines. Longitudinal, prospective data from the Whitehall study suggest that medically certified sickness absence can be used as a global measure of health. The question then arise whether gender differences merely is a mirror of women's higher burden of illness rather than based in social inequalities at work, or at home. Most studies on sickness absence control for health differences, but it is well known that women report higher levels of symptoms and poorer self-perceived health compared to men. In a longitudinal general population based study of women clinical psychiatric disorders were diagnosed at a personal interview. Other indicators of ill health in the study, were self-perceived mental and physical health. Women were then followed for five years and data on sickness absence were collected. At follow-up an analysis of health indicators as predictors of sickness absence over the five year period was performed. Since health problems differ between women and men, not only in occurrence but also in type of illnesses we choose a same sex data base for this analysis.

Lessons learned

Work environment is important in prevention of sickness absence in both men and women. However, sex differences might also be explained by differences in ill health. If so, the responsibility for the health care is higher to identify and treat women's health problems, and secondary prevention could be developed in order to avoid prolongation of sick-leave spells.

Conclusion

Differences in ill health can contribute to understand sex differences between women and men in sickness absence.

Abstract 2 title: Job and family related explanations of gender differences in sickness: A review of the evidence

Abstract 2 presenting authors: Arne Mastekaasa

Abstract 2 authors: Arne Mastekaasa

Affiliations 2: Department of Sociology, University of Oslo, Oslo, Norway

Abstract 2: Background

A prominent explanation of gender differences in sickness absence is that women to a greater extent than men have to combine employment with family or household related work and responsibilities. Another possible explanation is that women have less healthy jobs, or that they are more strongly affected by a given level of workplace exposures than are men. There is evidence for the importance of influence at work and exposure might differ between men and women. The gender segregated labour market has also been discussed as a possible factor contributing to the higher sickness absence in women, but findings are contradictory. This paper reviews the evidence with regard to these explanations. Methodological considerations related to selection bias in studies on sickness absence will be presented. Studies comparing men and women in different job and family situations as well as studies that more directly attempt to measure stressors at home and at work will be covered.

Abstract 3 title: Sickness absence as an interactive process: gendered experiences of young, highly educated women with mental health problems

Abstract 3 presenting authors: Petra Verdonk

Abstract 3 authors: Verdonk P 1, de Rijk A 1, Klinge I 2, de Vries A 3

Affiliations 3: 1 Department of Social Medicine, Maastricht University, Maastricht

2 Department of Health, Ethics & Society, Maastricht University, Maastricht

3 Beyaert foundation in Maastricht, the Netherlands

Abstract 3: Objective

Highly educated Dutch women experience more work related mental health disability than their male counterparts, and yet little is known regarding the process. Using the theory of symbolic interactionism, we examined how women interpret their roles at work, during sick leave, and upon their return to work.

Methods

Semi-structured interviews focusing on role perceptions and interactions with other actors were conducted with thirteen women (aged 29-41 years) on sick leave or off work for periods ranging from half a year to eight years.

Results

The women worked overtime because of workaholism, or to meet supervisors' expectations. This led to mental health problems and social isolation. Taking sick leave aided recovery, but further isolated the women. Insufficient support from the workplace and social insurance professionals intensified negative feelings. Psychological counselling provided alternatives whereby work and private roles could become more balanced. However, their reintegration into the workplace failed because the women could not implement these strategies when the organizational culture failed to change.

Conclusions

A long lead-up time preceded sickness absence and sick leave allowed for recovery and value adjustment. However, a variety of interpretations reinforced the women's individualized focus, thereby hampering their successful reintegration.

Practice implications

Given the importance of implementing effective sick leave prevention measures in the workplace, psychological treatment should focus on women's interactions with their work environment.

Abstract 4 title: Sickness absence in women – what are the associations to domestic work?

Abstract 4 presenting authors: Carin Staland Nyman

Abstract 4 authors: Staland Nyman C<sup>1</sup>, Alexanderson K<sup>2</sup>, Hensing G<sup>1</sup>

Affiliations 4: <sup>1</sup>Social Medicine, Department of Public Health and Community Medicine, The Sahlgrenska Academy at the University of Gothenburg

<sup>2</sup> Section of Personal Injury Prevention, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

Abstract 4: Issue

One explanation put forward for women's higher sickness absence, is they often have the main responsibility for the domestic work. Studies on the association between health and "double

burden” and “interactions between paid and unpaid work”, respectively are common. Few have addressed the association with sickness absence and the specific impact of domestic work.

#### Description of the problem

Compared to paid work, measures on exposure to domestic work are scarce and less developed. In previous studies on association between domestic work and sickness absence, different measures, such as number and age of children at home, responsibility for household work, working hours in domestic work, and interference between paid and unpaid work have been used. Results are inconsistent and improved measures have been asked for. In a study on employed women in Sweden a multidimensional perspective of domestic work was used in constructing different measures of domestic work. No associations were found between sickness absence and ‘Domestic job strain’. ‘Caring activities related to adults’, ‘Caring activities related to children’ and ‘Domestic life events or difficulties’ were associated with higher risk of sick-leave spells 8-30 days. Lacking ‘Domestic work equity and marital satisfaction’ was associated with a higher risk of a new sick-leave spell during the study period, while ‘Parental responsibility’ was associated with lower risk of sick-leave.

#### Lessons learned

Domestic work combines life domains and work tasks with very different meaning and content. A multidimensional assessment of domestic work contributed to identify specific aspects of domestic work that might affect women’s sickness absence, and that might be possible to prevent. Future research needs to improve measures focusing both general and specific domestic work aspects as well as contextual factors so that the complexity of domestic work becomes better defined and operationalised.